

Model structures on operads and algebras from a global perspective

David White

Denison University

Joint with Michael Batanin and Florian De Leger

IWoAT Conference for Peter May

March 3, 2025

arXiv:2311.07320 and arXiv:2311.07322

High level overview of my research

- Many of my papers **set up homotopy theory for operads and algebras** (model categories, localization).
- Method: work through a **complicated filtration** to understand cell attachments.
- Upshot: characterize when localization preserves algebraic structure (application: Kervaire paper), prove the existence of N_∞ -operads (Blumberg-Hill conjecture), prove the Baez-Dolan stabilization hypothesis.
- Flavors: Unital vs non-unital, reduced vs non-reduced, CF, nonsymmetric, cyclic, modular operads (Ginzberg and Kapranov), wheeled, n-operads, PROPs, properads...
- Approach as algebras over colored operads.

Overview of this talk

- ① Goal: unified approach to right induced model structures on operads, algebras, modules, bimodules. Here f a weak equivalence (resp. fibration) iff $U(f)$ is, where $U = \text{forgetful}$
- ② Different flavors of operads have different categories of trees, hence different filtrations.
- ③ First: extract filtration from category of trees in a categorical way.
- ④ Second: tweak trees to study operads and algebras/modules simultaneously. Get model structure on category of pairs (O, A) , or triples (O, A, M) or (O, P, A) , etc. Deduce model structures on operads, algebras, modules, etc. Recover known + new results.

A zoo of monoidal model categories

Throughout, let \mathcal{M} denote a nice monoidal model category, like:

- 1 Spaces: $(\text{sSet}, \times, *)$ or Top
- 2 Pointed spaces: $(\text{sSet}_*, \wedge, S^0)$ or Top_* . Equivariant, too.
- 3 Chain complexes: $(\text{Ch}(\mathbb{R}), \otimes_{\mathbb{R}}, S^0(\mathbb{R}))$
- 4 $(\mathbb{k}[G] - \text{mod}, \otimes_{\mathbb{k}}, \mathbb{k})$ with stable module category
- 5 (Categories of presheaves or functors, Day convolution)
- 6 Spectra, G -spectra, motivic spectra
- 7 Graphs, groupoids, categories, 2-cat, weak n -cat, etc.

Some of these have a structured interval object. Others require a hard filtration working cell-by-cell.

All of these satisfy our conditions for transferring model structures. **Do you have more model categories in mind?**

Grothendieck construction

- Given $\Phi : \mathcal{B}^{\text{op}} \rightarrow \text{CAT}$, form the Grothendieck construction $\int \Phi$ whose **objects are pairs** (O, A) where $O \in \mathcal{B}$ and $A \in \Phi(O)$, e.g., $O = \text{operad}$ and A is O -algebra.
- Morphisms $(\phi, f) : (O, A) \rightarrow (O', A')$ has $\phi : O \rightarrow O'$ and $f : A \rightarrow \phi^*(A')$.
- Global setting of $\int \Phi$ is a convenient place to study all ‘fibers’ (a.k.a. vertical structures) simultaneously.
- Examples: $\mathcal{B} = \text{monoids}$, and $\Phi(R) := R\text{-modules}$.
- $\mathcal{B} = \text{operads}$, and $\Phi(O) := \text{Alg}_O$ or $\Phi(O) := \text{Mod}_O$.
- $\mathcal{B} = \text{pairs of operads}$, and $\Phi(O, P) := (O, P)\text{-bimodules}$.
- $\mathcal{B} = (O, P, A, B, X)$, and A, B are (O, P) -bimod, and X infinitesimal (A, B) -bimod.
- **Goal:** connect homotopy theory of $\mathcal{B}, \Phi(O)$, and $\int \Phi$.

Related papers

- Previous papers assumed \mathcal{B} and all $\Phi(O)$ have model str's + **more**, then induced model str on $\int \Phi$, whose weak equivalences and fibrations match those in \mathcal{B} and $\Phi(O)$'s.
- Roig 1994, Stanculescu 2012. Assume:
 - 1 for every w.e. ϕ in \mathcal{B} , then ϕ^* preserves and reflects w.e.'s.
 - 2 for every triv cof u in \mathcal{B} , the unit of $(u_!, u^*)$ is a w.e.
- Harpaz-Prasma 2015:
 - 1 for every trivial cof. u or triv. fib. v in \mathcal{B} , then $u_!$ and v^* preserve weak equivalences.
 - 2 for every weak equiv. ϕ , $(\phi_!, \phi^*)$ is a Quillen equivalence.
- Cagne-Mellies 2020 (**conditions imply HP2**):
 - 1 HP1 but now $u_!$ and v^* preserve and reflect w.e.'s
 - 2 Beck-Chevalley. Given $u \circ v = v' \circ u$ in \mathcal{B} , then $\mu : (u')_! v^* \rightarrow (v')^* u_!$ is w.e. in $\Phi(\text{dom}(v'))$.

Our strategy is the opposite: get $\int \Phi$ first

- The assumptions in previous work often fail, e.g., the weak equivalence $O \rightarrow \text{Com}$ for an E_∞ -operad O does not induce a Quillen equivalence on algebras in spaces.
- For operad-style settings, **we induce a model structure on $\int \Phi$ from $\mathcal{M}^{\mathbb{N}+1}$, then deduce model structures on \mathcal{B} and all $\Phi(O)$** . Often $\int \Phi$ is alg over $\mathbb{N} + 1$ colored operad.
- **New filtration** for operads and algebras simultaneously.
Think: **trees with extra markings for algebras**.
- For operad-style settings, we always get a semi-model structure on $\int \Phi$ and hence induce same on \mathcal{B} and on $\Phi(O)$'s for cofibrant O .
- **Recover all known results** about (semi-)model structures on operads and algebras, **plus new results**, from one theorem.
- Key is polynomial monads and **quasi-tame** notion.

How to do homotopy theory in $\int \Phi$

Lemma (well-known)

If $T = UF$ is monad on cofibrantly gen. $\mathcal{N} = \text{Coll}(\mathcal{M}) \times \mathcal{M}$ and if for all generating trivial cofibrations $j : K \rightarrow L$ in \mathcal{N} , transfinite compositions of pushouts in $\text{Alg}_T(\mathcal{N})$:

$$\begin{array}{ccc} F(K) & \longrightarrow & F(L) \\ \downarrow & & \downarrow \\ (O, A) & \longrightarrow & (O', B) \end{array}$$

\Downarrow


are weak equivalences then $\text{Alg}_T(\mathcal{N})$ has **transferred model structure**, with weak equivalences and fibrations defined in \mathcal{N} .

If above works only for (O, A) cofibrant (resp. $U((O, A))$ cofibrant) then get semi-model structure (resp. semi over \mathcal{N}).

This cofibrancy means Σ_n -action on $O(n)$ is free; quotient easier.

Filtration to compute the pushout $(O, A) \rightarrow (O', B)$

New filtration covers operads and algebras at once via trees with marked (boxed) vertices plus rules for composition:



Now $(O, A) \rightarrow (O', B)$ is a transfinite composition of simpler pushouts in \mathcal{N} . Hence, $(O, A) \rightarrow (O', B)$ is a weak equivalence. If polynomial monad P is quasi-tame, we get a model structure on $\int \Phi$. Otherwise, we get a semi-model structure.

Semi-model categories, given $F : \mathcal{M} \rightleftarrows \mathcal{D} : U$

Definition: (\mathcal{D}, W, Q, F) satisfies all model category axioms except we only require the following for A and K cofibrant (resp. cof in \mathcal{M}):

$$\begin{array}{ccc} A & \longrightarrow & X \\ \cong \downarrow & \nearrow & \downarrow \\ B & \longrightarrow & Y \end{array} \quad \& \quad \begin{array}{ccc} K & \longrightarrow & L \\ \cong \searrow & & \nearrow \\ & D & \end{array}$$

Still have cofibrant replacement. **All model category results have semi-model category analogues** (often cofibrantly replace first): Ken Brown lemma, cylinders and path objects, cube lemma, Quillen equivalences, Reedy model structures, (co)simplicial frames, homotopy (co)limits, simplicial mapping spaces, Bousfield localization, etc. **Combinatorial semi is Quillen equiv. to combinatorial model category.**

Polynomial monads

A polynomial P from I to J in \mathbf{Set} is a diagram of sets of the form

$$I \xleftarrow{s} E \xrightarrow{p} B \xrightarrow{t} J$$

Such a diagram generates a functor:

$$\mathbb{P} : \mathbf{Set}/I \rightarrow \mathbf{Set}/J.$$

$$\mathbb{P}(X)_j = \coprod_{b \in t^{-1}(j)} \prod_{e \in p^{-1}(b)} X_{s(e)},$$

If $I = J$ and $p^{-1}(b)$ is finite for all b and P is a cartesian monad then $P\text{-alg}$ is equivalent to algebras over a Σ -cofibrant J -colored operad.

Polynomial monads encode flavors of operads

A finitary polynomial monad P has a category of algebras $\text{Alg}_P(\mathcal{M})$ (with I colors). **Plug in your flavor of trees.**

- **Free monoid monad.** $M(X) = \coprod_n X^n$. The corresponding polynomial is

$$1 \leftarrow \text{LTr}^* \rightarrow \text{LTr} \rightarrow 1$$

Where LTr^* is linear rooted trees with one vertex marked.

- **Free non symmetric operad monad.** The corresponding polynomial is

$$\mathbb{N} \leftarrow \text{PTr}^* \rightarrow \text{PTr} \rightarrow \mathbb{N}$$

Where PTr^* is planar rooted trees with one vertex marked.

- **Free symmetric operad monad.** The corresponding polynomial is

$$\mathbb{N} \leftarrow \text{ORTr}^* \rightarrow \text{ORTr} \rightarrow \mathbb{N}$$

Where ORTr is ordered rooted trees.

Examples of polynomial monads

- Free symmetric operad monad; also non-symmetric operads; presheaves, monoids, enriched categories;
- Monads for cyclic and modular operads;
- Dioperads, properads, (generalized) PROPs, and wheeled and colored versions of all monads above.
- Free n -operad monad (see Batanin-Berger ‘Tame’ paper).
- New: categories of pairs (O, A) , triples (O, A, M) , (O, P, A) , (O, P, A, B, M) , etc.

So, all these are also algebras over appropriate Σ -cofibrant colored operads.

Polynomial monads and the Grothendieck construction

- **Goal:** If \mathcal{B} is the category of algebras over a polynomial monad, then so is $\int \Phi$.
- Let T be an I -colored symmetric operad in \mathcal{M} , equipped with a morphism of operads $\phi : T \rightarrow \text{SO}(J)$. Note: ϕ induces $\phi^* : \text{Alg}_{\text{SO}_p(J)}(\mathcal{M}) \rightarrow \text{Alg}_T(\mathcal{M})$.
- Let O be an algebra of T . An algebra of O in \mathcal{M} is a J -collection $C = \{C_j \mid j \in J\}$ of objects of \mathcal{M} equipped with a map of T -algebras $O \rightarrow \phi^*(\text{End}(C))$ where $\text{End}(C)$ is the endomorphism operad of C .
- The category of O -algebras is isomorphic to the category $\Phi(O) := \text{Alg}_{\phi_!(O)}(\mathcal{M})$.
- **Theorem:** If ϕ is a map of polynomial monads in Set then there exists a polynomial monad $\text{Gr}(T)$ such that the category $\int \Phi$ is isomorphic to the category $\text{Alg}_{\text{Gr}(T)}(\mathcal{M})$.

Polynomial monad $\text{Gr}(T)$

We are given $\phi : T \rightarrow \text{SO}(J)$, displayed vertically:

$$\begin{array}{ccccccc}
 I & \xleftarrow{s} & E & \xrightarrow{p} & B & \xrightarrow{t} & I \\
 \downarrow c & & \downarrow & & \downarrow \psi & & \downarrow c \\
 \text{Bq}(J) & \xleftarrow{\quad} & \text{ORTr}^*(J) & \xrightarrow{\quad} & \text{ORTr}(J) & \xrightarrow{\quad} & \text{Bq}(J)
 \end{array}$$

We define a polynomial monad

$$I \sqcup J \xleftarrow{S} D^* \xrightarrow{\quad} D \xrightarrow{T} I \sqcup J$$

where $D = B \sqcup B = B \sqcup \{(b, \sigma) \mid \sigma \in \text{ORTr}(J), b \in \psi^{-1}(\sigma)\}$ and S and T are induced by s, t, c, ψ above.

Upshot: If T is a polynomial monad then so is $\text{Gr}(T)$, hence we have a semi-model structure on $\int \Phi$ always.

Analyze the pushout P of $X \xleftarrow{g} F(K) \xrightarrow{F(f)} F(L)$

- Construct a monad $T_{f,g}$ whose algebras are 5-tuples (X, K, L, f, g) . There's a map of monads $a : T_{f,g} \rightarrow T$ s.t. $a_! : \text{Alg}_{T_{f,g}}(\mathcal{M}) \rightarrow \text{Alg}_T(\mathcal{M})$ is exactly the pushout P .
- $U(P)$ is the **colimit over the classifier $T^{T_{f,g}}$, a categorified simplicial bar construction** that picks out $T_{f,g}$ -algebras in T -algebras. This $T^{T_{f,g}}$ decomposes like words in X, K, L . So $U(X) \rightarrow U(P)$ is a transfinite composition of pushouts of morphisms in \mathcal{M} .
- T is **tame if $T^{T_{f,g}}$ has a discrete final subcategory**; colimit easy to compute. Get full model structure on Alg_T .
- T is **quasi-tame if $\pi_1(T^{T+1})$ is equivalent to a discrete groupoid**. Get full model structure. Tame implies quasi-tame. Monad T for nonsymmetric operads is tame but $\text{Gr}(T)$ is only quasi-tame.

Model category conditions on monoidal model cat \mathcal{M}

- **Monoid axiom:** (Triv. Cof. $\otimes \mathcal{M}$)-cell \subset w.e.'s.
- **h-monoidal:** for each (trivial) cofibration $f : X \rightarrow Y$ and each object Z , the map $f \otimes Z$ is a (trivial) h-cofibration, i.e.,

$$\begin{array}{ccccc} X \otimes Z & \longrightarrow & A & \xrightarrow{\simeq} & B \\ \downarrow f \otimes Z & & \downarrow & & \downarrow \\ Y \otimes Z & \longrightarrow & A' & \xrightarrow{\simeq} & B' \end{array}$$

- **Compactly generated:** all objects are small relative to I^\otimes -cell, and weak equivalences are closed under filtered colimits along morphisms in I^\otimes .
- **Commutative monoid axiom:** if f is a trivial cof then so is $f^{\square n} / \Sigma_n$. Ex: \mathbf{sSet} , \mathbf{Top} , $\mathbf{Ch}(k)$, $\mathbf{StMod}(k[G])$, $\mathbf{spectra}^+$.
Yields (\mathbf{R}, \mathbf{M}) model str for \mathbf{R} commutative monoid.

We get model structures on:

- Pairs (R, M) where R is a monoid and M is an R -module.
- Pairs (O, A) where O is a nonsymmetric operad and A is an O -algebra. Same for (O, M) with left O -module.
- Triples (O, P, M) where M is an (O, P) -bimodule. **Infinitesimal, too**, via (O, P, A, B, M) .
- Pairs (O, M) where O is a constant-free symmetric operad (or n -operad) and M is a constant-free module.
- Semi on (O, A) where O is a symmetric operad (or hyperoperad, modular, cyclic, PROP, dioperad, properad, wheeled, etc) and A is an O -algebra.
- **Let $\mathcal{M} = \text{Ch}(k)$, k characteristic zero. Get full (vertical) model structure on category of twisted modular operads of Ginzberg and Kapranov (1998). This is new.**

Structure of Grothendieck construction

- Assume everything is bicomplete (need to make them model categories) and all ϕ^* preserve w.e.'s.
- Projection $p : \int \Phi \rightarrow \mathcal{B}$, $p(O, A) = O$.
- Right adjoint $r : \mathcal{B} \rightarrow \int \Phi$, $r(O) = (O, t_O)$.
- Left adjoint $i : \mathcal{B} \rightarrow \int \Phi$, $i(O) = (O, i_O)$.
- Sometimes $E : \int \Phi \rightarrow \mathcal{B}$ like enveloping algebra or enveloping operad $(O, A) \mapsto O_A$.

Global to horizontal and vertical

- All (semi-)model structures are transferred, so we know the weak equivalences and fibrations in $\mathcal{B}, \Phi(\mathcal{O}), \int \Phi$.
- Note (ϕ, f) is a w.e. (resp. fib) iff ϕ and f are. Define cofibrations by the lifting property.
- Lemma: $(\phi, f) : (\mathcal{O}, A) \rightarrow (\mathcal{P}, B)$ is a **global (trivial) cofibration** if and only if ϕ is a horizontal (trivial) cofibration and $f^* : \phi_!(A) \rightarrow B$ is a vertical (trivial) cofibration.
- **Theorem:** If $\int \Phi$ admits global (semi-)model structure then \mathcal{B} admits horizontal (semi-)model structure, $\Phi(\mathcal{O})$ admits vertical model structure for each $\mathcal{O} \in \mathcal{B}$ (semi-model structure for a cofibrant \mathcal{O}), and $p : \int \Phi \rightarrow \mathcal{B}$ is left and right Quillen.
- If $\phi : \mathcal{O} \rightarrow \mathcal{O}'$, then ϕ^* and $\phi_!$ form a Quillen pair.

Cofibrant generation, properness, and rectification

- If $\int \Phi$ is **left (resp. right) proper** then \mathcal{B} and $\Phi(O)$ are left (right) proper for any O . Same for **cofibrantly generated**.
- Same for relatively left/right proper.
- **Application:** for $\mathcal{M} = \text{Ch}(k)$, characteristic zero, then $O\text{-alg}$ is **left proper** for all O .
- If $\int \Phi$ is left proper, and ϕ is w.e., ϕ^* reflects w.e.'s, and the unique map $\tau : i_O \rightarrow \phi^*(i_{O'})$ is a w.e., then **$(\phi^*, \phi_!)$ is a Quillen equivalence** (this is rectification). Application: **strictification in Lack's model structure** for 2Cat .
- Relative version if relatively left proper and O, O' are u-cofibrant, for $(U, u) : \int \Phi \rightarrow \int \Psi$. Like Σ -cofibrant.
- If O, O' are cofibrant then get Q.E. even if $\int \Phi$ is not (relatively) left proper.

From semi to full

Lemma

If \mathcal{M} is a semi-model and any morphism admits (triv. cof, fib.) factorization then \mathcal{M} is a model category.

So: lifting follows from factorization.

Lemma

Let $\int \Phi$ be semi and all ϕ^* preserve fibrations and weak equivalences. If, for any $f : A \rightarrow B$ in $\Phi(\mathcal{O})$, the induced map $E(\mathcal{O}, f) : E(\mathcal{O}, A) \rightarrow E(\mathcal{O}, B)$ admits a (triv. cof., fib.) factorization in \mathcal{B} , then so does f in $\Phi(\mathcal{O})$.

Upshot: if \mathcal{B} is full model structure then so are the $\Phi(\mathcal{O})$'s.

Future Work

- De Leger: hyperoperads, triple delooping, infinitesimal bimodules.
- Bousfield localization for $\int \Phi$.
- For n-operad global model structure, force quasi-bijections act invertibly. Prove global Baez-Dolan stabilization result.
- Generalize Braun, Chuang, Lazarev work on derived localizations of (A, M) where A is an algebra and M is an A -module, to global setting $\int \Phi$, e.g., (O, A) where O is operad and A is O -algebra.
- Relate localizations of operads and algebras.

References

- Batanin-Berger: Homotopy theory for algebras over polynomial monads, TAC.
- Batanin-De Leger: Polynomial monads and delooping of mapping spaces, Noncomm. Geom. 2020.
- Batanin-De Leger-White: Model structures on operads and algebras from a global perspective, arXiv:2311.07320.
- Batanin-De Leger-White: Quasi-tame substitutes and the Grothendieck construction, arXiv:2311.07322.
- Batanin-White: Homotopy theory of algebras of substitutes and their localisation, Transactions AMS.
- Cagne-Melliès: On bifibrations of model categories, Advances.
- De Leger: Cofinal morphism of polynomial monads and double delooping, arXiv:2205.09149.
- De Leger-Grego: Triple delooping for multiplicative hyperoperads, arXiv:2309.15055, and new work on Lattice Path Operad.
- Harpaz-Prasma: The Grothendieck construction for model categories.
- Roig: Model category structures in bifibred categories, JPAA.
- Stanculescu: Bifibrations and weak factorisation systems, ACS.
- White-Yau: Bousfield localization and algebras over colored operads.